

**AETNA BETTER HEALTH® OF VIRGINIA REQUEST FORM**

**MULTIPLE SCLEROSIS**

**Fax back to: 1-855-799-2553**

If the following information is not complete, correct, or legible, the PA process can be delayed. Please use one form per member.

**MEMBER INFORMATION**

**Last Name:**

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

**First Name:**

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

**Medicaid ID Number:**

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

**Date of Birth:**

				-					-										
--	--	--	--	---	--	--	--	--	---	--	--	--	--	--	--	--	--	--	--

**Weight in Kilograms:** \_\_\_\_\_

**PRESCRIBER INFORMATION**

**Last Name:**

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

**First Name:**

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

**NPI Number:**

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

**Phone Number:**

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

**Fax Number:**

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

**DRUG INFORMATION**

**Drug Name/Form:** \_\_\_\_\_

**Strength:** \_\_\_\_\_

**Dosing Frequency:** \_\_\_\_\_

**Length of Therapy:** \_\_\_\_\_

**Quantity per Day:** \_\_\_\_\_

*(Form continued on next page.)*

**Member's Last Name:**

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

**Member's First Name:**

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

**DIAGNOSIS AND MEDICAL INFORMATION**

---

1. Is the member at least 18 years of age?

Yes     No

2. Has the member had a baseline magnetic resonance imaging (MRI) before initiating the first treatment course (within 3 months prior to start of therapy)?

Yes     No

3. Indicate all that apply:

Relapsing-remitting disease (RRMS)       Secondary progressive disease (SPMS) with relapses

Clinically isolated syndrome (CIS)       Member has had ≥ 1 relapse within the previous two years

Member has new and unequivocally enlarging T2 contrast enhancing lesions as evidenced by MRI and has had ≥ 1 relapse in the previous 12 months

Other: \_\_\_\_\_

4. Has the member had a treatment failure or contraindication to other agents used to treat multiple sclerosis (MS)? List previous medications (include drug name/dose):

Yes     No

Previous Medication(s): \_\_\_\_\_

5. Will Mavenclad®, Mayzent®, Ponvory™, Zeposia® be used as single-agent therapy?

Yes     No

6. Has the member been tested for antibodies to the varicella zoster virus (VZV) or received immunization for VZV four weeks prior to beginning therapy?

Yes     No

7. Has the member been screened for the presence of tuberculosis according to local guidelines?

Yes     No

8. Has the member been evaluated and screened for the presence of hepatitis B and hepatitis C virus (HBV/HCV) prior to initiating treatment?

Yes     No

*(Form continued on next page.)*

**Member's Last Name:**

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

**Member's First Name:**

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

**9. Mavenclad® Specific**

a. Is the lymphocyte count  $\geq$  800 cells/mL prior to start of therapy?

Yes     No

b. Please attest that members of childbearing age are not pregnant **and** that members of reproductive potential must use effective contraception during treatment with therapy and for at least six months after the last dose.

Yes     No

c. Does the member have human immunodeficiency virus (HIV) infection?

Yes     No

**10. Mayzent® Specific**

a. Has the member been tested for CYP2C9 variant status to determine genotyping (required for dosing)?

Yes     No

**11. Mayzent®, Ponvory™ or Zeposia® Specific**

a. Please attest that members of childbearing age are not pregnant and that members of reproductive potential must use effective contraception during treatment.

Yes     No

b. Has the member obtained a baseline electrocardiogram (ECG)?

Yes     No

c. Has the member had a baseline ophthalmic evaluation of the fundus, including the macula, before starting treatment?

Yes     No

**12. Before using Mayzent®, Ponvory™ or Zeposia®, can you attest that the member does not have any of the following:**

- Recent myocardial infarction
- Unstable angina
- Stroke
- Transient ischemic attack
- Decompensated heart failure with hospitalization
- Class III/IV heart failure within the previous 6 months
- Prolonged QTc interval at baseline (> 500 msec)
- CYP2C9\*3/\*3 genotype (**Mayzent® only**)
- History of Mobitz Type II second or third-degree atrioventricular block or sick sinus syndrome (unless treated with a functioning pacemaker)

Yes     No

*(Form continued on next page.)*

Member's Last Name:

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

Member's First Name:

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

13. Can you confirm that **Mayzent®** will **not** be used in combination with the following?:

- Moderate or strong CYP3A4 inducers (e.g., modafinil, efavirenz) in members with a CYP2C9\*1/\*3 and CYP2C9\*2/\*3 genotypes; **OR**
- Drug regimens that contain CYP2C9/CY3A4 dual inhibitors (e.g., fluconazole); **OR**
- Moderate CYP2C9 inhibitor plus a moderate-to-strong CYP3A4 inhibitor; **OR**
- Other antineoplastic, immunosuppressive or immunomodulating drugs.

Yes     No

14. Can you confirm **Zeposia®** will **not** be used in combination with the following?:

- Will **not** be initiating therapy after previous treatment with alemtuzumab; **OR**
- Monoamine oxidase inhibitor (MAOI) (e.g., selegiline, phenelzine, linezolid); **OR**
- Drugs known to prolong the QT-interval (e.g., fluoroquinolone or macrolide antibiotics, venlafaxine, fluoxetine, quetiapine, ziprasidone, sumatriptan, zolmitriptan); **OR**
- Strong cytochrome p450 2C8 (CYP2C8) inhibitors (e.g., gemfibrozil) or inducers (e.g., rifampin); **OR**
- BCRP inhibitors (e.g., cyclosporine, eltrombopag); **OR**
- Adrenergic or serotonergic drugs which can increase norepinephrine or serotonin (e.g., opioids, selective serotonin reuptake inhibitors [SSRIs], selective norepinephrine reuptake inhibitors [SNRIs], tricyclics, tyramine); **OR**
- Foods with large amounts of tyramine (e.g., > 150 mg), such as aged cheeses, cured meats, craft/unfiltered beers, beans); **OR**
- Other antineoplastic, immunosuppressive or immunomodulating drugs (**Note:** if there is a history of prior use of these drugs, consider possible unintended additive immunosuppressive effects); **AND**
- Patient will **not** receive live vaccines during and at least 4 weeks prior to and 12 weeks after treatment; **AND**
- Patient does **not** have an active infection, including clinically important localized infections

Yes     No

---

Prescriber Signature (Required)

Date

By signature, the physician confirms the above information is accurate and verifiable by member records.

**Please include ALL requested information; Incomplete forms will delay the PA process.**

Submission of documentation does NOT guarantee coverage.