



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Acthar-Purified Corticotropin Gel Page: 1 of 4

Effective Date: 3/13/2025 Last Review Date: 2/2025

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> New Jersey
	<input checked="" type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Florida Kids	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Arizona

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Acthar and Purified Corticotropin Gel under the patient's prescription drug benefit.

Description:

Indications

The Limited Indication Specialty Guideline Management (LI SGM) program provides coverage for specific, but not all FDA labeled or compendial supported drug uses based on plan design and the scope of the pharmacy benefit. This program provides coverage for Acthar Gel for the treatment of infantile spasms and exacerbations of multiple sclerosis and coverage for Purified Cortrophin Gel for the treatment of exacerbations of multiple sclerosis if all of the approval criteria are met.

- A. **Infantile Spasms (Acthar Gel only):** as monotherapy for the treatment of infantile spasms in infants and children under 2 years of age
- B. **Multiple Sclerosis:** treatment of acute exacerbations of multiple sclerosis in adults

The use of Acthar and Purified Cortrophin Gel for the treatment of all other indications listed in the FDA product labeling has not been proven to be superior to conventional therapies (e.g., corticosteroids, immunosuppressive agents) and has a significantly higher cost than the standard of care agents. Use of Acthar and Purified Cortrophin Gel for these conditions is considered not medically necessary and is not a covered benefit.

A. Acthar Gel:

1. **Rheumatic Disorders:** as adjunctive therapy for short-term administration (to tide the patient over an acute episode or exacerbation) in: psoriatic arthritis; rheumatoid arthritis, including juvenile rheumatoid arthritis; ankylosing spondylitis
2. **Collagen Diseases:** during an exacerbation or as maintenance therapy in selected cases of: systemic lupus erythematosus, systemic dermatomyositis (polymyositis)
3. **Dermatologic Diseases:** severe erythema multiforme, Stevens-Johnson syndrome
4. **Allergic States:** serum sickness
5. **Ophthalmic Diseases:** severe acute and chronic allergic and inflammatory processes involving the eye and its adnexa such as: keratitis, iritis, iridocyclitis, diffuse posterior uveitis and choroiditis, optic neuritis, chorioretinitis, anterior segment inflammation
6. **Respiratory Diseases:** symptomatic sarcoidosis
7. **Edematous State:** to induce a diuresis or a remission of proteinuria in nephrotic syndrome without uremia of the idiopathic type or that due to lupus erythematosus



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name:	Acthar-Purified Corticotropin Gel	Page:	2 of 4
Effective Date:	3/13/2025	Last Review Date:	2/2025
Applies to:	<input checked="" type="checkbox"/> Illinois <input checked="" type="checkbox"/> Maryland <input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Florida <input checked="" type="checkbox"/> Florida Kids <input type="checkbox"/> Virginia	<input checked="" type="checkbox"/> New Jersey <input type="checkbox"/> Michigan <input type="checkbox"/> Arizona

B. Purified Cortrophin Gel:

- Rheumatic Disorders:** as adjunctive therapy for short-term administration (to tide the patient over an acute episode or exacerbation) in: psoriatic arthritis; rheumatoid arthritis, including juvenile rheumatoid arthritis; ankylosing spondylitis; acute gouty arthritis.
- Collagen Diseases:** during an exacerbation or as maintenance therapy in selected cases of: systemic lupus erythematosus, systemic dermatomyositis (polymyositis).
- Dermatologic Diseases:** severe erythema multiforme (Stevens-Johnson syndrome), severe psoriasis
- Allergic States:** atopic dermatitis, serum sickness
- Ophthalmic Diseases:** severe acute and chronic allergic and inflammatory processes involving the eye and its adnexa such as: allergic conjunctivitis, keratitis, iritis and iridocyclitis, diffuse posterior uveitis and choroiditis, optic neuritis, chorioretinitis, anterior segment inflammation
- Respiratory Diseases:** symptomatic sarcoidosis
- Edematous States:** to induce a diuresis or a remission of proteinuria in the nephrotic syndrome without uremia of the idiopathic type or that due to lupus erythematosus

Note: Acthar Gel single-dose pre-filled SelfJect injector is not to be used for the treatment of infantile spasms.

All other indications are considered experimental/investigational and not medically necessary.

Applicable Drug List:

Acthar gel (repository corticotropin injection)
Purified Cortrophin gel (repository corticotropin injection)

Policy/Guideline:

Documentation:

Submission of the following information is necessary to initiate the prior authorization review for requests for treatment of multiple sclerosis exacerbations: chart notes detailing the outcome of the most recent trial with IV methylprednisolone, including dosage and duration of treatment.

Exclusions:

- A. Coverage of Purified Cortrophin Gel for the treatment of infantile spasms will be excluded.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Acthar-Purified Corticotropin Gel Page: 3 of 4
Effective Date: 3/13/2025 Last Review Date: 2/2025

Applies to: Illinois Florida New Jersey
 Maryland Florida Kids Michigan
 Pennsylvania Kids Virginia Arizona

B. Use of Acthar Gel in combination with Purified Cortrophin Gel will be excluded.

Criteria for Initial Approval:

A. Infantile Spasms (Acthar Gel only)

Authorization of 4 weeks may be granted for treatment of infantile spasms in members who are less than 2 years of age.

B. Multiple Sclerosis

Authorization of 3 weeks may be granted for treatment of acute exacerbations of multiple sclerosis when the member has had an inadequate response to a trial of IV methylprednisolone (for the current exacerbation).

Continuation of Therapy:

A. Infantile Spasms (Acthar Gel only)

Authorization of 4 weeks may be granted to members requesting Acthar Gel for continuation of therapy when the member has shown substantial clinical benefit from therapy.

B. Multiple Sclerosis

Authorization of 3 weeks may be granted for members requesting re-authorization of repository corticotropin therapy when all initial authorization criteria are met.

Approval Duration and Quantity Restrictions:

Approval:

- Infantile spasms – 4 weeks
- Multiple Sclerosis – 3 weeks

Quantity Level Limits:

- Acthar Gel mutli-dose vial 80 units/mL for IM or SC injection – 35 mL per 21 days
- Acthar Gel 40 units/0.5 mL single-dose pre-filled SelfJect injector for SC injection – 28 injectors per 28 days
- Acthar Gel 80 units/0.5 mL single-dose pre-filled SelfJect injector for SC injection – 28 injectors per 28 days
- Purified Corticotropin Gel multi-dose vial 80 units/mL for IM or SC injection – 35 mL per 21 days

References:

1. Acthar Gel [package insert]. Bedminster, NJ: Mallinckrodt ARD LLC.; February 2024.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Acthar-Purified Corticotropin Gel Page: 4 of 4

Effective Date: 3/13/2025 Last Review Date: 2/2025

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input checked="" type="checkbox"/> New Jersey
	<input checked="" type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Florida Kids	<input type="checkbox"/> Michigan
	<input checked="" type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Arizona

2. Pellock JM, Hrachovy R, Shinnar S, et al. Infantile spasms: A U.S. consensus report. *Epilepsia*. 2010;51:2175-2189.
3. Go CY, Mackay MT, Weiss SK, et al. Evidence-based guideline update: Medical treatment of infantile spasms: Report of the Guideline Development Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology*. 2012;78:1974-1980.
4. Hancock EC, Osborne JP, Edwards SW. Treatment of infantile spasms. *Cochrane Database Syst Rev*. 2013;6:CD001770.
5. Riikonen R. Recent advances in pharmacotherapy of infantile spasms. *CNS Drugs* 2014; 28:279-290.
6. Pavone P, et al. Infantile spasms syndrome, West Syndrome and related phenotypes: what we know in 2013. *Brain & Development* 2014; 739-751.
7. Citterio A, La Mantia L, Ciucci G, et al. Corticosteroids or ACTH for acute exacerbations in multiple sclerosis. *Cochrane Database Syst Rev* 4:CD001331.
8. Thompson AJ, Kennard C, Swash M, et al. Relative efficacy of intravenous methylprednisolone and ACTH in the treatment of acute relapse in MS. *Neurology* 1989; 39:969-971.
9. Berkovich R, Agius MA. Mechanisms of action of ACTH in the management of relapsing forms of multiple sclerosis. *Ther Adv Neurol Disord* 2014; 7(2):83-96.
10. Frohman EM, Shah A, Eggenberger E, et al. Corticosteroids for multiple sclerosis: I. Application for treatment exacerbations. *Neurotherapeutics* 2007; 4(4): 618-626.
11. Sellebjerg F, Barnes D, Filippini G, et al. EFNS guidelines on treatment of multiple sclerosis relapses: report of an EFNS task force on treatment of multiple sclerosis. *European Journal of Neurology* 2005; 12:939-946.
12. Purified Cortrophin Gel [package insert]. Baudette, MN: ANI Pharmaceuticals, Inc.; October 2023.