



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Nucala

Page: 1 of 6

Effective Date: 2/26/2025

Last Review Date: 1/29/2025

Applies to: New Jersey

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Nucala under the patient's prescription drug benefit.

Description:

- A. Nucala is indicated for add-on maintenance treatment of patients with severe asthma aged 6 years and older, and with an eosinophilic phenotype (*Reference Biologics in Severe Asthma New Jersey State Protocol.*)
- B. Nucala is indicated for the treatment of adult patients with eosinophilic granulomatosis with polyangiitis (EGPA).
- C. Nucala is indicated for the treatment of adult and pediatric patients aged 12 years and older with hypereosinophilic syndrome (HES) for ≥ 6 months without an identifiable non-hematologic secondary cause.
- D. Nucala is indicated for add-on maintenance treatment of adult patients 18 years and older with chronic rhinosinusitis with nasal polyps (CRSwNP).

If the member is a current smoker or vaper, they should be counseled on the harmful effects of smoking and vaping on pulmonary conditions and available smoking and vaping cessation options.

All other indications are considered experimental/investigational and not medically necessary.

For all indications:

Member will not use the requested medication concomitantly with any other biologic drug or targeted synthetic drug for the same indication.

Applicable Drug List:

Nucala

Policy/Guideline:

Criteria for Initial Approval:

Submission of the following information is necessary to initiate the prior authorization review:

- 1. Eosinophilic granulomatosis with polyangiitis**
 - i. Member's chart or medical record showing pretreatment blood eosinophil count.
- 2. Hypereosinophilic syndrome (HES)**



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Nucala

Page:

2 of 6

Effective Date: 2/26/2025

Last Review Date: 1/29/2025

Applies to: New Jersey

- i. FIP1L1-PDGFR α fusion gene test results.
- ii. Member's chart or medical record showing pretreatment blood eosinophil count.

3. Chronic rhinosinusitis with nasal polyps

- i. Member's chart or medical record showing nasal endoscopy, anterior rhinoscopy, or computed tomography details (e.g., location, size), or Meltzer Clinical Score or endoscopic nasal polyps score (NPS) (where applicable).
- ii. Chart notes, medical record documentation, or claims history supporting previous medications tried. If therapy is not advisable, documentation of clinical reason to avoid therapy.

Nucala must be prescribed by or in consultation with ONE of the following:

- i. Chronic rhinosinusitis with nasal polyposis: allergist/immunologist or otolaryngologist

Initial Criteria:

Authorization may be granted for treatment of asthma when ALL the following criteria are met:

1. Eosinophilic granulomatosis with polyangiitis (EGPA)

- i. Member is 18 years of age or older.
- ii. Member has a history or the presence of an eosinophil count of more than 1000 cells per microliter or a blood eosinophil level of greater than 10%.
- iii. Member is currently taking oral corticosteroids, unless contraindicated or not tolerated
- iv. Member has at least TWO of the following disease characteristics of EGPA:
 - a. Biopsy showing histopathological evidence of eosinophilic vasculitis, perivascular eosinophilic infiltration, or eosinophil-rich granulomatous inflammation
 - b. Neuropathy, mono or poly (motor deficit or nerve conduction abnormality)
 - c. Pulmonary infiltrates, non-fixed; sino-nasal abnormality
 - d. Cardiomyopathy (established by echocardiography or magnetic resonance imaging)
 - e. Glomerulonephritis (hematuria, red cell casts, proteinuria)
 - f. Alveolar hemorrhage (by bronchoalveolar lavage)
 - g. Palpable purpura
 - h. Anti-neutrophil cytoplasmic anti-body (ANCA) positive (Myeloperoxidase or proteinase 3)



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Nucala

Page:

3 of 6

Effective Date: 2/26/2025

Last Review Date: 1/29/2025

Applies to: New Jersey

- v. Member has had at least one relapse (requiring increase in oral corticosteroids dose, initiation/increased dose of immunosuppressive therapy or hospitalization) within 2 years prior to starting treatment with Nucala or has a refractory disease

2. Hypereosinophilic syndrome (HES)

- i. Member is 12 years of age or older.
- ii. Member does not have EITHER of the following:
 - a. HES secondary to a non-hematologic cause (e.g., drug hypersensitivity, parasitic helminth infection, [human immunodeficiency virus] HIV infection, non-hematologic malignancy)
 - b. FIP1L1-PDGFR α kinase-positive HES
- iii. Member has a history or presence of a blood eosinophil count of at least 1000 cells per microliter.
- iv. Member will not use Nucala as monotherapy.
- v. Member has been on a stable dose of HES therapy (e.g., oral corticosteroid, immunosuppressive, and/or cytotoxic therapy).
- vi. Member has had HES for at least 6 months.
- vii. Member has experienced at least two HES flares within the past 12 months

3. Chronic rhinosinusitis with nasal polyps

- i. Member is 18 years of age or older
AND
Members has previously received a biologic drug indicated for Chronic rhinosinusitis with nasal polyps (CRSwNP) in the past year.
 - a. Requests will require that the patient is unable to take Dupixent and Xolair for the given diagnosis due to a trial and inadequate treatment response or intolerance, or a contraindication. Documentation is required for approval.
- OR
- ii. Member has bilateral nasal polyposis and chronic symptoms of sinusitis despite intranasal corticosteroid treatment for at least 2 months unless contraindicated or not tolerated; and
- iii. The member has CRSwNP despite ONE of the following:
 - a. Prior sino-nasal surgery; or
 - b. Prior treatment with systemic corticosteroids within the last two years was ineffective, unless contraindicated or not tolerated
- iv. Member has ONE of the following:



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Nucala

Page:

4 of 6

Effective Date: 2/26/2025

Last Review Date: 1/29/2025

Applies to: New Jersey

- a. A bilateral nasal endoscopy, anterior rhinoscopy, or computed tomography (CT) showing polyps reaching below the lower border of the middle turbinate or beyond in each nostril
- b. Meltzer Clinical Score of 2 or higher in both nostrils
- c. A total endoscopic nasal polyp score (NPS) of at least 5 with a minimum score of 2 for each nostril
- v. Member has nasal blockage plus ONE additional symptom:
 - a. Rhinorrhea (anterior/posterior); or
 - b. Reduction or loss of smell; or
 - c. Facial pain or pressure
- vi. Member will continue to use a daily intranasal corticosteroid while being treated with Nucala, unless contraindicated or not tolerated.

Criteria for Continuation of Therapy:

1. Eosinophilic granulomatosis with polyangiitis (EGPA)

- i. Chart notes or medical record documentation supporting improvement in EGPA control
- ii. Member is 18 years of age or older.
- iii. Member has beneficial response to treatment with Nucala as demonstrated by any of the following:
 - a. A reduction in the frequency of relapses, or
 - b. A reduction or discontinuance of daily oral corticosteroid dose, or
 - c. No active vasculitis

2. Hypereosinophilic syndrome (HES)

- i. Chart notes or medical record documentation supporting improvement in HES control.
- ii. Member is 12 years of age or older.
- iii. Member has experienced a reduction in HES flares since starting treatment with Nucala.
- iv. Member will not use Nucala as monotherapy.

3. Chronic rhinosinusitis with nasal polyps

- i. Chart notes or medical record documentation supporting positive clinical response.
- ii. Member is 18 years of age or older.
- iii. Member has achieved or maintained a positive clinical response to Nucala therapy as evidenced by improvement in signs and symptoms of CRSwNP (e.g., improvement in nasal congestion, nasal polyp size, loss of smell,



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name:	Nucala	Page:	5 of 6
Effective Date:	2/26/2025	Last Review Date:	1/29/2025
Applies to:	<input checked="" type="checkbox"/> New Jersey		

anterior or posterior rhinorrhea, sinonasal inflammation, hyposmia and/or facial pressure or pain or reduction in corticosteroid use).

- iv. Member will continue to use a daily intranasal corticosteroid while being treated with the requested medication, unless contraindicated or not tolerated.

Approval Duration and Quantity Restrictions:

Initial Approval

- Chronic rhinosinusitis with nasal polyps: 6 months
- Eosinophilic granulomatosis with polyangiitis: 12 months
- Hypereosinophilic syndrome (HES): 12 months

Renewal Approval: 12 months

Quantity Level Limit:

- Nucala 100 mg single-dose vial: 3 vials per 28 days
- Nucala 100 mg/mL single-dose prefilled safety syringe: 3 syringes per 28 days
- Nucala 100 mg/mL single-dose prefilled autoinjector: 3 autoinjector's per 28 days
- Nucala 40mg/0.4mL, single-dose prefilled syringe: 1 syringe per 28 days

References:

1. Nucala [package insert]. Durham, NC: GlaxoSmithKline; March 2023.
2. Ortega HG, Liu MC, Pavord ID, et al. Mepolizumab treatment in patients with severe eosinophilic asthma. *N Engl J Med.* 2014;371(13):1198-1207.
3. Bel EH, Wenzel SE, Thompson PJ, et al. Oral glucocorticoid-sparing effect of mepolizumab in eosinophilic asthma. *N Engl J Med.* 2014;371(13):1189-1197.
4. National Institutes of Health. National Asthma Education and Prevention Program Expert Panel Report 3: Asthma Management Guidelines: Focused Updates 2020. Bethesda, MD: National Heart Lung and Blood Institute; December 2020. Available at: <https://www.nhlbi.nih.gov/sites/default/files/publications/AsthmaManagementGuidelinesReport-2-4-21.pdf>. Accessed March 12, 2024.
5. Global Initiative for Asthma (GINA). Global Strategy for Asthma Management and Prevention. 2023 update. Available at: https://ginasthma.org/wp-content/uploads/2023/07/GINA-Full-Report-23_07_06-WMS.pdf. Accessed March 12, 2024.
6. Kew KM, Karner C, Mindus SM. Combination formoterol and budesonide as maintenance and reliever therapy versus combination inhaler maintenance for chronic asthma in adults and children (review). *Cochrane Database Syst Rev.* 2013;12:CD009019.
7. American Academy of Allergy, Asthma & Immunology (AAAAI) 2020 Virtual Annual Meeting. Available at: <https://annualmeeting.aaaai.org/>. Accessed March 12, 2024.
8. [Wechsler ME](#), [Akuthota P](#), [Jayne D](#), et al. Mepolizumab or placebo for eosinophilic granulomatosis with polyangiitis. *N Engl J Med.* 2017;18;376(20):1921-1932.
9. GlaxoSmithKline. A Study to Investigate Mepolizumab in the Treatment of Eosinophilic Granulomatosis With Polyangiitis. Available from <https://clinicaltrials.gov/ct2/show/record/NCT02020889>. NLM identifier: NCT02020889. Accessed March 14, 2024.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name:	Nucala	Page:	6 of 6
Effective Date:	2/26/2025	Last Review Date:	1/29/2025
Applies to:	<input checked="" type="checkbox"/> New Jersey		

10. Groh M, Pagnoux C, Baldini C, et al. Eosinophilic granulomatosis with polyangiitis (Churg–Strauss) (EGPA) Consensus Task Force Recommendations for evaluation and management. *Eur J Intern Med.* 2015;26(7):545-553.
11. Yates M, Watts RA, Bajema M, et al. EULAR/ERA-EDTA recommendations for the management of ANCA-associated vasculitis. *Ann Rheum Dis.* 2016;75(9):1583-1594.
12. Shomali W, Gotlib J. World Health Organization-defined eosinophilic disorders: 2022 update on diagnosis, risk stratification, and management. *Am J Hematol.* 2022;97(1):129-148.
13. Butt NM, Lambert J, Ali S, et al. Guideline for the investigation and management of eosinophilia. *Br J Haematol.* 2017;176(4):553-572.
14. Han JK, Bachert C, Fokkens W, et al. Mepolizumab for chronic rhinosinusitis with nasal polyps (SYNAPSE): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Respir Med.* 2021;9(10):1141-1153.
15. Bachert C, Han JK, Wagenmann M, et al. EUFOREA expert board meeting on uncontrolled severe chronic rhinosinusitis with nasal polyps (CRSwNP) and biologics: Definitions and management. *J Allergy Clin Immunol.* 2021;147(1):29-36.
16. Cloutier MM, Dixon AE, Krishnan JA, et al. Managing asthma in adolescents and adults: 2020 asthma guideline update from the National Asthma Education and Prevention Program. *JAMA.* 2020;324(22):2301-2317.
17. American College of Rheumatology. 2021 American college of rheumatology/vasculitis foundation guideline for the management of antineutrophil cytoplasmic antibody-associated vasculitis. *Arthritis & Rheumatology.* <https://www.vasculitisfoundation.org/wp-content/uploads/2021/07/2021-ACR-VF-Guideline-for-Management-of-ANCA-Associated-Vasculitis.pdf>. Accessed March 14, 2024.
18. Fokkens WJ, Lund VJ, Hopkins C, et al. European Position Paper on Rhinosinusitis and Nasal Polyps 2020. *Rhinology.* 2020;58(Suppl S29):1-464.
19. Hopkins C. Chronic Rhinosinusitis with Nasal Polyps. *N Engl J Med.* 2019;381(1):55-63